

## Birth Control Use After Delivery

Percentage (%) of recent mothers in Illinois who reported using the following birth control methods

Among women reporting using birth control now

Survey Question 64: What kind of birth control are you or your husband or partner using now to keep from getting pregnant? Response: Yes

	Percentage <sup>1</sup>	95% CI <sup>2</sup>		Respondents <sup>3</sup>	Estimated Population Affected <sup>4</sup>
Condoms	29.2	26.1	32.5	950	29,703
Birth control pills	23.1	20.2	26.2	950	23,503
IUD (including Mirena®, ParaGard®, Liletta®, or Skyla®)	16.6	14.1	19.4	950	16,857
Withdrawal (pulling out)	14.4	12.0	17.1	950	14,613
Not having sex (abstinence)	9.3	7.5	11.6	950	9,489
Tubes tied or blocked (female sterilization or Essure®)	8.0	6.3	10.2	950	8,184
Shots or injections (Depo-Provera®)	7.1	5.4	9.2	950	7,187
Contraceptive implant in the arm (Nexplanon® or Implanon®)	5.7	4.2	7.6	950	5,771
Natural family planning (including rhythm method)	4.0	2.9	5.6	950	4,060
Vasectomy (male sterilization)	3.9	2.8	5.5	950	3,985
Contraceptive patch (OrthoEvra®) or vaginal ring (NuvaRing®)	2.3	1.4	3.7	950	2,367

<sup>1</sup>Percentage = Weighted percentage<sup>2</sup>CI = Confidence interval lower/upper bounds

-- Indicates denominator &lt;30 respondents or numerator &lt;6 respondents

<sup>3</sup>Respondents = The total number who responded to the survey question<sup>4</sup>Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition