
Prescription Pain Reliever Education

Percentage (%) of recent mothers in Illinois who reported a health care worker discussed affects on baby of using prescription pain relievers during pregnancy

Survey Question 012: At any time during your most recent pregnancy, did a doctor, nurse, or other health care worker talk with you about how using prescription pain relievers during pregnancy could effect a baby? Response: Yes



		Percentage ¹	95% Cl ²		Respondents ³	Estimated Population Affected4
					•	
All	Overall	52.9	49.3	56.5	898	50,743
Maternal Age	<20 yrs	63.2	45.0	78.3	35	2,467
	20 to 24 yrs	67.9	58.5	76.0	137	12,728
	25 to 29 yrs	50.5	43.7	57.4	242	12402.4
	30 to 34 yrs	46.9	40.7	53.1	285	13,660
	=>35 yrs	48.4	40.8	56.0	199	9,485
Race/Ethnicity	NH White	44.6	39.8	49.6	468	22,684
	NH Black	75.5	66.7	82.6	148	10977.5
	Hispanic	54.4	46.8	61.9	200	12,140
	Other	61.5	48.9	72.7	76	4630.0
Education	<12 yrs	73.0	62.0	81.7	92	6864.9
	12 yrs	61.0	53.2	68.4	199	13,450
	>12 yrs	47.5	43.2	51.9	601	30,392
Locale	Urban	52.8	48.7	56.8	725	40,579
	Rural	53.3	45.1	61.4	173	10,163
Birth Weight	Low	53.0	46.7	59.3	240	3,794
	Normal	52.9	49.0	56.7	658	46,949
Insurer for Delivery	Medicaid	65.5	59.6	70.9	339	23,786
	Other	44.7	40.3	49.3	554	26,342
WIC	Yes	68.7	62.1	74.6	250	18,977
	No	46.4	42.1	50.7	640	31,296

¹Percentage = Weighted percentage

Data collection period April - December 2019

 $Other\ Insurer\ for\ Delivery = Self-pay,\ Indian\ Health\ Service,\ Champus/Tricare,\ Other\ Government,\ Other\ Government$

²CI = Confidence interval lower/upper bounds

⁻⁻ Indicates denominator <30 or numerator <6

³Respondents = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition