

## Illinois PRAMS FACTS – 2018

Pregnancy Risk Assessment Monitoring System



The Illinois Pregnancy Risk Assessment Monitoring System (PRAMS) is an ongoing population-based survey of women who delivered live born infants in Illinois. The Illinois PRAMS project is part of a national initiative of the U.S. Centers for Disease Control and Prevention (CDC) to reduce infant morbidity and mortality. More Illinois PRAMS information and detailed data can be found at <a href="http://www.dph.illinois.gov/data-statistics/prams">http://www.dph.illinois.gov/data-statistics/prams</a>.

SELECTED INDICATORS	PERCENTAGE% <sup>1</sup>	95% Cl <sup>2</sup>	Respondents <sup>3</sup>	Estimated Population Affected <sup>4</sup>		
Before Pregnancy						
During the 12 months before pregnancy						
<ul> <li>Exercised three or more days per week</li> </ul>	41.8	38.9-44.7	541	57,040		
Dieted to lose weight	30.6	27.9-33.4	378	41,637		
<ul> <li>Had teeth cleaned by a dentist or dental hygienist</li> </ul>	58.6	55.2-62.0	563	58,438		
<ul> <li>Had a health care visit with an OB/GYN</li> </ul>	63.1	59.7-66.4	595	62,881		
Had health care visit for illness or chronic condition	15.6	13.3-18.3	152	15,564		
Had health care visit for family planning or birth control	12.5	10.4-14.9	121	12,448		
Had health care visit for depression or anxiety	9.5	7.7-11.8	91	9,520		
Checked for diabetes by health care worker	20.3	18.0-22.8	268	27,560		
Taking regular prescription medications, other than birth control	21.2	18.9-23.8	285	28,988		
Talked with a health care worker about their family history	42.8	39.9-45.7	565	58,333		
During the 3 months before pregnancy						
<ul> <li>Diagnosed with diabetes</li> </ul>	2.3	1.6-3.4	34	3,145		
Diagnosed with high blood pressure	4.3	3.3-5.7	64	5,918		
Diagnosed with depression	12.3	10.5-14.4	159	16,834		
Took multivitamins, prenatal vitamins, or folic acid vitamins every day during the month before pregnancy	38.5	35.7-41.4	508	52,507		
Pregnancy unintended – want to be pregnant later, not at all, or not sure	42.1	39.2-45.1	544	57,266		
Smoked during the three months before pregnancy	14.7	12.7-17.0	196	19,794		
Consumed alcoholic drinks during the three months before pregnancy	63.0	60.1-65.8	802	85,929		
Were physically abused by current or ex-husbands/partners, a family member, or someone else during 12 months before pregnancy	2.9	2.1-4.1	40	3,941		
Enrolled in Medicaid the month before pregnancy	30.1	27.4-32.9	400	40,302		

SELECTED INDICATORS	PERCENTAGE% <sup>1</sup>	95% Cl <sup>2</sup>	Respondents <sup>3</sup>	Estimated Population Affected <sup>4</sup>			
During Pregnancy							
Received prenatal care during the first trimester	88.5	86.4-90.3	1,136	119,742			
Used WIC services during pregnancy	28.7	26.1-31.5	374	38,451			
Had teeth cleaned by a dentist or dental hygienist	42.8	39.9-45.8	557	58,233			
Diagnosed with depression	11.1	9.3-13.1	148	15,056			
Received influenza immunization 12 months before or during pregnancy	61.1	58.2-64.0	778	82,611			
Diagnosed with high blood pressure (starting in pregnancy) or (pre-)eclampsia	15.5	13.5-17.7	244	20,955			
Diagnosed with gestational diabetes	9.3	7.7-11.2	124	12,659			
Were physically abused by current or ex-husbands/partners, a family	2.6	1.8-3.7	32	3,448			
member, or someone else during pregnancy							
Smoked last three months of pregnancy	6.9	5.5-8.5	93	9,243			
Felt unsafe in the neighborhood where living	8.5	6.9-10.3	114	11,568			
Enrolled in Medicaid for prenatal care	43.7	40.7-46.7	556	57,411			
Postpartum Experiences and Infant Health							
Smoked after delivery	9.6	7.9-11.5	128	12,895			
Diagnosed with depression	11.2	9.5-13.3	148	15,238			
Had a postpartum checkup	93.0	91.3-94.4	1,202	126,645			
Used postpartum birth control	80.2	77.8-82.5	1,023	109,098			
Breastfed at least once after birth	87.0	84.8-88.8	1,102	117,013			
Breastfed for at least 12 weeks after birth	57.1	54.1-60.1	678	73,571			
Infant usual sleeping							
<ul> <li>Infant put to sleep on their backs most of the time</li> </ul>	84.2	81.8-86.2	1,066	113,204			
<ul> <li>Infant slept in crib/bassinet/play yard</li> </ul>	90.8	88.9-92.4	1,155	121,380			
Infant slept with a blanket (not recommended)	46.1	43.1-49.1	566	60,294			
Enrolled in Medicaid after delivery	37.1	34.2-40.0	473	49,599			

<sup>&</sup>lt;sup>1</sup>Percentage = Weighted percentage of Illinois mothers who gave birth to live infants in 2018 and responded Yes or as indicated to the question for the specified indicator.

The 2018 Illinois PRAMS survey was sent to 2,175 recent mothers in Illinois who delivered live born infants. The number of mothers responding to the survey was 1,306 for an unweighted response rate of 60%. The overall weighted response rate was 61%. The weighted response rates were 56% for mothers of low birthweight infants and 62% for mothers of normal birthweight infants. (Source: Illinois 2018 Analysis Weights report.)

<sup>&</sup>lt;sup>2</sup>CI = Confidence Interval

<sup>&</sup>lt;sup>3</sup>Respondents (N) = Total number of mothers who responded Yes to the survey question for the specified indicator; <30 respondents results may be unreliable <sup>4</sup>Estimated population affected = Estimated number of Illinois mothers who delivered live born infants in 2018 and who would respond Yes or as indicated to the question for the specified indicator.