## Birth Control Method Used After Delivery<sup>1</sup> Among Women Who Reported Current Use of Birth Control Illinois PRAMS 2012

	Percentage <sup>2</sup>	95% Confidence Interval	Respondents <sup>3</sup>	Estimated Population Affected <sup>4</sup>
Condoms	31.3	28.1 - 34.8	280	38060
Birth control pills	22.9	19.9 - 26.1	201	27790
IUD (including Mirena® or ParaGard®)	14.0	11.6 - 16.8	111	16960
Withdrawal (pulling out)	12.4	10.2 - 15.0	104	15077
Tubes tied or blocked (female sterilization)	6.6	5.0 - 8.7	58	8060
Injections (Depo-Provera®)	6.1	4.6 - 8.2	63	7427
Natural family planning (including rhythm method)	4.9	3.6 - 6.6	44	5896
Not having sex (abstinence)	4.6	3.3 - 6.4	43	5630
Contraceptive implant (Implanon®)	3.8	2.6 - 5.6	28	4641
Vasectomy (male sterilization)	3.5	2.4 - 5.1	29	4289
Contraceptive patch (OrthoEvra®) or vaginal ring (NuvaRing®)	2.6	1.7 - 4.1	24	3177

<sup>&</sup>lt;sup>1</sup>Survey Question 68: What kind of birth control are you or your husband or partner using now to keep from getting pregnant?

<sup>&</sup>lt;sup>2</sup>**Percentage** = Weighted percentage of Illinois mothers giving birth in 2012 with the specified indicator for each characteristic.

<sup>&</sup>lt;sup>3</sup>Respondents (N) = Total number of mothers who responded to this question with Yes; total N = 1075 (Yes or No); Total N value for each characteristic varies due to

<sup>&</sup>lt;sup>4</sup>Estimated population affected = Estimated number of Illinois mothers giving birth in 2012 with the specified indicator for each characteristic