Before & During Pregnancy

Table 37

Other Tobacco Products

Percentage (%) of recent mothers in Illinois who reported using other tobacco products in the past two years

Survey Question 37: Have you used any of the following products in the past 2 years? Response: Yes

	Percentage ¹	95% Cl ²		Respondents ³	Estimated Population Affected ⁴
E-cigarettes or other electronic nicotine products	6.0	4.7	7.6	1,293	8,099
Hookah	4.2	3.2	5.6	1,289	5,684
Cigars, cigarillos, or little filered cigars	2.1	1.4	3.1	1,289	2,798
Chewing tobacco, snuff, snus, or dip	0.5	0.2	1.1	1,288	678

Percentage (%) of recent mothers in Illinois who reported using other tobacco products in the three months before pregnancy or the last three months of pregnancy

Among women who reported using other tobacco products in the past two years

Survey Question 38: During the 3 months before you got pregnant, on average, how often did you use e-cigarettes or other electornic nicotine products? Responses 'Yes/No' created using numeric response

Survey Question 39: During the last 3 months of your pregnancy, on average, how often did you use e-cigarettes or other electronic nicotine products? Responses 'Yes/No' created using numeric response

	Percentage ¹	95% Cl ²		Respondents ³	Estimated Population Affected ⁴
E-cigarette use 3 months before pregnancy	3.5	2.5	4.8	1,296	4,715
E-cigarette use last 3 months of pregnancy	1.3	0.8	2.2	1,297	1,756

¹Percentage = Weighted percentage

²**CI** = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents or numerator <6 respondents

³**Respondents** = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition