Birth Control Use After Delivery

Percentage (%) of recent mothers in Illinois who reported using the following birth control methods Among women reporting using birth control now

Survey Question 64: What kind of birth control are you or your husband or partner using now to keep from getting pregnant? Response: Yes

	Percentage ¹ 95% Cl ²		6 CI ²	Respondents ³	Estimated Population Affected4
Condoms	31.4	28.5	34.5	1,111	37,311
Birth control pills	22.7	20.1	25.6	1,111	26,974
IUD (including Mirena®, ParaGard®, Liletta®, or Skyla®)	16.0	13.7	18.6	1,111	18,994
Withdrawal (pulling out)	13.7	11.6	16.1	1,111	16,290
Tubes tied or blocked (female sterilization or Essure®)	7.8	6.2	9.8	1,111	9,232
Shots or injections (Depo-Provera®)	7.6	6.1	9.6	1,111	9,082
Not having sex (abstinence)	7.2	5.8	9.1	1,111	8,600
Natural family planning (including rhythm method)	5.1	3.8	6.6	1,111	6,016
Contraceptive implant in the arm (Nexplanon® or Implanon®)	3.9	2.8	5.5	1,111	4,633
Vasectomy (male sterilization)	3.1	2.2	4.5	1,111	3,729
Contraceptive patch (OrthoEvra®) or vaginal ring (NuvaRing®)	2.2	1.4	3.5	1,111	2,638

¹Percentage = Weighted percentage

²CI = confidence interval

⁻⁻ Indicates denominator <30 respondents

³Respondents = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition