After Delivery

Table 57


2016 Phase 8 Illinois PRAMS Data

Birth Control Use After Delivery

Percentage (%) of recent mothers in Illinois who who reported using a most or moderately effective* form of birth control

Among women who were not trying to get pregnant and who were not currently pregnant

Survey Question 64: What kind of birth control are you or your husband or partner using now to keep from getting pregnant? Response: Yes to a most or moderately effecitve method



		Percentage ¹	95% Cl ²		Respondents ³	Estimated Population Affected ⁴
All	Overall	54.2	51.2	57.2	1,275	74,282
Maternal Age	<20 yrs	77.1	61.9	87.4	51	4,779
	20 to 24 yrs	61.0	53.2	68.2	205	14,965
	25 to 29 yrs	57.5	51.5	63.2	334	22,437
	30 to 34 yrs	50.4	45.3	55.6	425	21,063
	=>35 yrs	43.5	37.0	50.2	260	11,039
Race/Ethnicity	NH White	55.2	51.3	59.1	712	42,377
	NH Black	56.1	47.8	64.1	197	10,240
	Hispanic	59.9	52.7	66.7	237	17,755
	Other	30.4	21.6	40.9	109	3167.4
Education	<12 yrs	64.7	55.5	73.0	148	11,301
	12 yrs	60.0	53.0	66.5	250	17,384
	>12 yrs	50.4	46.7	54.1	843	43,813
Birth Weight	Low	54.0	48.6	59.2	338	5,156
	Normal	54.3	51.0	57.5	937	69,126
Marital Status	Married	48.7	45.0	52.5	791	41,022
	Other	63.0	58.0	67.8	484	33,260
Insurer Now	Medicaid	67.2	62.3	71.8	475	34,919
	Other	46.3	42.6	50.1	800	39,363

¹Percentage = Weighted percentage

²**CI** = confidence interval

-- Indicates denominator <30 respondents

³**Respondents** = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Most Moderately Effective Method includes male/female sterilization, birth control pills, implant (e.g., Nexplanon, Implanon*), IUD (e.g., Mirena*, ParaGard*, Liletta*, Skyla*), contraceptive patch (e.g. OrthoEvra*) or shot (Depo-Provera*), or vaginal ring (e.g., NuvaRing*).