
Birth Control Use After Delivery

Percentage (%) of recent mothers in Illinois who who reported using a most or moderately effective* form of birth control

Among women who were not trying to get pregnant and who were not currently pregnant

Survey Question 64: What kind of birth control are you or your husband or partner using now to keep from getting pregnant? Response: Yes to a most or moderately effecitve method



All	Overall	Percentage ¹ 50.9	95% Cl ²		Respondents ³	Estimated Population Affected ⁴
			47.7	54.1	1,159	68,448
Maternal Age	<20 yrs	66.2	48.1	80.6	41	3,008
	20 to 24 yrs	55.3	47.2	63.0	194	12,927
	25 to 29 yrs	46.5	40.3	52.7	304	16,837
	30 to 34 yrs	48.7	43.2	54.2	377	20,427
	=>35 yrs	53.9	47.0	60.6	243	15,248
Race/Ethnicity	NH White	51.2	46.9	55.5	600	36,706
	NH Black	47.1	39.2	55.2	215	10,187
	Hispanic	60.3	53.2	67.1	232	17,323
	Other	30.3	21.4	41.1	99	3238.6
Education	<12 yrs	57.1	47.3	66.4	136	9,471
	12 yrs	56.8	49.4	64.0	226	15,907
	>12 yrs	47.7	43.9	51.5	771	41,455
Birth Weight	Low	52.1	46.7	57.5	324	5,176
	Normal	50.8	47.4	54.2	835	63,271
Marital Status	Married	46.5	42.7	50.5	713	36,746
	Other	57.1	51.8	62.3	446	31,701
Insurer Now	Medicaid	55.5	50.0	60.8	436	28,032
	Other	48.2	44.2	52.1	723	40,415

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

⁻⁻ Indicates denominator <30 respondents

³Respondents = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

^{*}Most Moderately Effective Method includes male/female sterilization, birth control pills, implant (e.g., Nexplanon*, Implanon*), IUD (e.g., Mirena*, ParaGard*, Liletta*, Skyla*), contraceptive patch (e.g. OrthoEvra*) or shot (Depo-Provera*), or vaginal ring (e.g., NuvaRing*).