




# CHLAMYDIA IS THE MOST COMMONLY REPORTED DISEASE IN ILLINOIS

Every year there are around 65,000 cases of chlamydia in Illinois. Chlamydia is transmitted through unprotected anal, oral and vaginal sex. Untreated chlamydia infections can lead to complications like pelvic inflammatory disease (PID), ectopic pregnancies and infertility. Women are at a greater risk than men of getting a chlamydia infection.

## Who has chlamydia?



7 of 10 cases are in women



Have no symptoms

**Simple Testing** performed on urine or swabs

**2 Pills** standard treatment



Use a condom

Have sex with only one uninfected partner (who only has sex with you)


Abstain from sexual activity

**GET TESTED AND REDUCE YOUR RISK!**


# GONORRHEA CAN HAVE SERIOUS SIDE EFFECTS IF LEFT UNTREATED

Every year there are around 17,000 cases of gonorrhea in Illinois. Gonorrhea can cause an infection in the anus, eyes, mouth, penis, throat, urinary tract or uterus and is transmitted through unprotected anal, oral and vaginal sex. Many women and men with gonorrhea do not have symptoms.


## Who has gonorrhea?



Cases are equal in men and women



## Cases are:



## Untreated Gonorrhea

- Increases risk of getting HIV
- Can cause infertility
- Can spread to blood, heart, skin and joints

**Simple Testing** performed on urine or swabs

**2 Drugs** standard treatment



Use a condom

Have sex with only one uninfected partner (who only has sex with you)



Abstain from sexual activity

**GET TESTED AND REDUCE YOUR RISK!**

# SYPHILIS MOST COMMONLY REPORTED IN MEN

Every year there are around 1,600 cases of syphilis in Illinois. Syphilis is transmitted through direct contact with syphilis sores during unprotected anal, oral and vaginal sex. Early symptoms of syphilis are a sore and later symptoms may be a rash, headache, fever, sore throat, aches, hair loss, and swollen joints. If syphilis is not treated, it can cause arthritis, blindness, heart disease, mental illness, and death.

## Who has syphilis?



### HIV Co-infection

- Sores make it easier to get HIV during sex
- About 35% of people with syphilis also are infected with HIV

**Simple Testing** One blood draw

**Simple Treatment**  
One dose of medicine for most cases

Use a condom

Have sex with only one uninfected partner (who only has sex with you)

Abstain from sexual activity

**GET TESTED AND REDUCE YOUR RISK!**